

* Author preprint. The final publication is available at www.springerlink.com.

Kunda, M. & Goel, A. (In press). "Thinking in Pictures as a cognitive account of autism." *Journal of Autism and Developmental Disorders*. Published online: 20 November 2010. Available at: <http://www.springerlink.com/content/x33g41837j317352/>

Thinking in Pictures as a Cognitive Account of Autism

Maithilee Kunda and Ashok K. Goel
Design & Intelligence Lab, School of Interactive Computing
Georgia Institute of Technology
85 Fifth Street NW, Atlanta, GA 30332, USA

Abstract

We analyze the hypothesis that some individuals on the autism spectrum may use visual mental representations and processes to perform certain tasks that typically developing individuals perform verbally. We present a framework for interpreting empirical evidence related to this "Thinking in Pictures" hypothesis and then provide comprehensive reviews of data from several different cognitive tasks, including the n -back task, serial recall, dual task studies, Raven's Progressive Matrices, semantic processing, false belief tasks, visual search, spatial recall, and visual recall. We also discuss the relationships between the Thinking in Pictures hypothesis and other cognitive theories of autism including Mindblindness, Executive Dysfunction, Weak Central Coherence, and Enhanced Perceptual Functioning.

Key words: Autism; cognition; information processing; mental imagery; verbal representations; visual representations; visual reasoning.

Corresponding author: mkunda@gatech.edu

Introduction

Numerous individuals on the autism spectrum have posited that they tend to use visual mental representations instead of verbal ones (e.g. Hurlburt et al. 1994). In her well-known autobiographical book *Thinking in Pictures*, for example, Temple Grandin (2006) describes how her visual thinking style benefits her work in engineering design but also creates difficulties in understanding abstract concepts. Among cognitive theorists in the autism research community, this “Thinking in Pictures” idea seems to have received limited focused and sustained consideration. This relative lack of attention perhaps is due not only to the introspective nature of the above accounts but also because the hypothesis seems ill-defined.

The purpose of this article is to refine one formulation of the Thinking in Pictures (TiP) hypothesis about cognition in autism and examine existing empirical evidence relating to this hypothesis, expanding on our previous work (Kunda & Goel, 2008). Our formulation of this hypothesis has two main parts:

- Assumption: Typically developing (TD) individuals are, in general, able to use both visual and verbal mental representations.
- Hypothesis: A subset of individuals on the autism spectrum exhibits a disposition towards using visual mental representations (and a corresponding bias against using verbal mental representations).

For the remainder of this paper, this (and only this) is what we mean by the TiP hypothesis.

Although for a time cognitive science debated whether visual mental representations even existed, the weight of evidence now seems to indicate that they do; they are usually described as being analogical (i.e. having some structural correspondence to what they represent) and closely tied to perceptual mechanisms (Kosslyn et al. 2006). In contrast, verbal mental representations are often described as being propositional (Pylyshyn, 2002). However, our reading of the literature on cognition in autism indicates that, like the literature on cognitive science in general, different interpretations of visual and verbal representations are often used in practice, usually on a task-by-task basis. In this paper, we will pin down precise meanings for *visual* and *verbal* in our discussions of individual tasks.

General evidence suggesting a visual/verbal disparity among individuals on the autism spectrum can be found in studies of cognitive profiles, or patterns of verbal (V) versus nonverbal (NV) intelligence as measured by standardized IQ tests. Some studies have noted a $V < NV$ (lower verbal than nonverbal IQ) pattern among individuals on the autism spectrum (Lincoln et al. 1988), though such findings have not been universal (Klin et al. 1995; Siegel et al. 1996). Joseph et al. (2002) found that, while children with autism were generally more likely to have a V-NV discrepancy in either direction than were TD children, children with autism having a $V < NV$ pattern of abilities showed greater social impairment than the other children with autism, irrespective of absolute levels of verbal or general ability. The distinctiveness of the $V < NV$ profile, and also its association with variables of diagnostic interest, led the authors to conjecture that such a profile might indicate “an etiologically significant subtype of autism” reflecting fundamental changes in cognition and neuroanatomy, rather than just the selective sparing of certain nonverbal abilities.

A tendency to exhibit a $V < NV$ profile is exactly what one might expect from an individual who thinks in pictures, with one important caveat: the standard tasks used to measure verbal and nonverbal abilities in IQ tests have been selected through extensive study of

neurotypical development and performance, and there is no guarantee that a test measuring a particular cognitive ability in TD individuals measures the same cognitive ability in individuals with autism, as there may be multiple different strategies that can be used to solve the same task. (We return to this point in the following section.)

Other behavioral data from autism suggesting an over-reliance on visual representations span many different cognitive and task domains (e.g. Heaton et al. 2008; Joseph et al. 2005; Whitehouse et al. 2006). On the neurobiological side, Mottron et al. (2006) reported that, across a variety of fMRI studies, individuals with autism tend to show increased brain activation in posterior, visual-perceptual brain regions and decreased activation in frontal brain regions often used for verbal processing.

For the remainder of this article, we first describe what sorts of predictions about behavior can be made using the TiP hypothesis, and then we give several examples of relevant empirical data from behavior and neurobiology. We conclude by discussing the relationship of the TiP hypothesis with several existing cognitive theories of autism, including Mindblindness, Executive Dysfunction, Weak Central Coherence, and Enhanced Perceptual Functioning.

Effects of Thinking in Pictures on Behavior

A simplistic consideration of the TiP hypothesis might lead to predictions that individuals with autism will show good performance on visual tasks and poor performance on verbal tasks. However, there are two different ways to classify tasks as *visual* or *verbal*: how a task **can** be solved (i.e. what sorts of mental representations and inferences are sufficient, but not necessary) and how tasks are **typically** solved (i.e. what sorts of mental representations and inferences do TD individuals generally use).




Figure 1. Task classifications according to how they **can** be solved (solid and dashed circles) and how they are **typically** solved (light and dark grey shadings).

Figure 1 illustrates the potential overlap between these two types of task classifications. The solid and dashed circles (A and B) represent tasks that **can** be solved visually or verbally, respectively, and their intersection ($A \cap B$) represents tasks that can be solved either way. For example, matching one of two very similar shades of red to a target red patch can be solved using visual representations but not using verbal ones (at least not easily), and so this task lies

inside solid circle **A** but outside dashed circle **B**. On the other hand, determining which of the words *shoe* or *now* rhymes with the word *too* can be solved using phonological verbal representations but not using visual ones, and so this task lies inside dashed circle **B** but outside solid circle **A**. Finally, deciding which of two red and green colored patches matches a target red patch can be solved using either visual or verbal representations (e.g. by matching on visual hue or on linguistic label), and so this task lies in the intersection $\mathbf{A} \cap \mathbf{B}$.

The light grey and dark grey shaded regions (\mathbf{T}_A and \mathbf{T}_B) represent tasks that are **typically** solved visually or verbally, respectively. The bulk of psychological evidence on how most humans solve cognitive tasks has given us \mathbf{T}_A and \mathbf{T}_B , by definition, and it is tempting to treat these classifications as the final answer on whether a task is visual or verbal. However, for a typically verbal task in \mathbf{T}_B , if that task happens to also be solvable visually (i.e. lies within $\mathbf{A} \cap \mathbf{B}$), it is possible that an individual disinclined to use verbal representations can use a compensatory visual strategy to successfully solve that task.

By making these distinctions, the performance of an individual on a given task (e.g. level of success) can be evaluated independently of their strategy selection (e.g. visual or verbal). Keeping this in mind, we now use the TiP hypothesis to make general predictions about the behavior of individuals with autism on three different types of tasks, as shown in Table 1.

Table 1: General Behavioral Predictions for Autism from the TiP Hypothesis

Prediction	Task type in Fig. 1	Task type description	TD strategy	TD performance	AU strategy	AU performance
P1	exclusively in B	tasks that can only be done verbally	verbal	successful	visual	impaired
P2	in \mathbf{T}_A	tasks typically done visually	visual	successful	visual	successful
P3	in $\mathbf{T}_B \cap \mathbf{A}$	tasks typically done verbally that can be done visually	verbal	successful	visual	successful

Note. AU = Individuals with autism; TD = Typically developing individuals

The first prediction is, perhaps, the least useful for testing the TiP hypothesis, as impaired performance on verbal-only tasks is unlikely to inform us about what mental representations an individual who thinks in pictures is using; for instance, such individuals may not be engaging any task-relevant representations at all. Also, data from these tasks will not be very useful as a point of distinction between the TiP hypothesis and other deficit accounts of autism. However, this TiP prediction is consistent with general evidence for verbal impairments in autism (*DSM-IV-TR*, 2000), though more precise relationships remain to be determined.

Regarding the second prediction, that individuals with autism use visual strategies to solve tasks that are also typically solved visually, a conservative claim might be that the visual strategies used by the two groups are the same, and therefore no behavioral differences in either task performance or strategy selection ought to be observed. However, there is significant evidence for behavioral differences in autism on typically visual tasks, ranging from changes in low-level perception (e.g. Bertone et al. 2005) to superior performance on certain visual tasks like the Embedded Figures Task (e.g. Jolliffe & Baron-Cohen, 1997). One possible TiP explanation of these differences is that a bias towards using visual representations leads to a general “visual expertise” not shared by TD individuals. However, these findings can also be

interpreted as indications of other forms of atypical cognitive processing, for example of greater detail-oriented processing (Happé & Frith, 2006) or superior low-level perceptual abilities (Mottron et al. 2006). If such processing differences are an integral aspect of autism, one important question for TiP will be how such differences might be related to a visual representation bias. In general, data from typically visual tasks are not necessarily the best test of the TiP hypothesis, as they alone cannot distinguish between the TiP account and other cognitive theories that posit superior visual processing in autism.

The third prediction, regarding tasks typically solved verbally that can also be solved visually, is the most useful for directly testing the TiP hypothesis. In particular, for a given task in this category, it should be possible to design experiments that illuminate whether the underlying representational strategy used by an individual is visual or verbal. Furthermore, experiments testing this third TiP prediction will provide the surest means for distinguishing TiP from other cognitive theories of autism, as (insofar as we have seen) no other cognitive account explicitly posits visual/verbal representational differences.

In the following two sections, we review empirical data related to the third and second TiP predictions, respectively. In particular, we look at:

- Tasks typically done verbally that can be done visually: (1) the *n*-back task, (2) serial recall, (3) dual task studies, (4) Raven's Progressive Matrices, (5) semantic processing, and (6) false belief tasks.
- Tasks typically done visually: (7) visual search, (8) spatial recall, and (9) visual recall.

TiP Prediction #3: Tasks Typically Done Verbally That Can Be Done Visually

The n-Back Task

In the *n*-back task (Kirchner, 1958), a subject is presented with a sequence of stimuli and asked whether the current stimulus matches the one shown *n* steps ago. The variable *n* can take the value of one (respond "yes" to any succession of two identical stimuli), two (respond "yes" to any stimulus matching the one presented two steps back), and so on. Stimuli can vary as to their content and presentation, such as letters presented visually or auditorily, pictures, etc.

For TD individuals, the *n*-back task is thought to recruit verbal rehearsal processes in working memory (i.e. *phonological* verbal representations), among other executive resources (Smith & Jonides, 1999). Several published studies of the *n*-back task have not shown significant differences in accuracy or reaction time for individuals with autism relative to TD controls,¹ which has led, in some cases, to the conclusion that verbal working memory is intact in autism (Williams et al. 2005).

However, recent fMRI studies have shown that, while behavioral measures on the *n*-back task may be similar, there can be significant differences in patterns of brain activation between individuals with autism and TD controls. In one study using stimuli of visually presented letters, the autism group showed less brain activation than controls in left prefrontal and parietal regions associated with verbal processing and greater activation in right hemisphere and posterior

¹ See Appendix A.

regions associated with visual processing (Koshino et al. 2005). In another study using stimuli of photographs of faces, a similar decrease in left prefrontal activation was found in the autism group (Koshino et al. 2008). Both of these studies suggest that individuals with autism may be using a visual strategy for the *n*-back task, whereas controls use at least a partially verbal strategy.

Serial Recall

In serial recall tasks, a subject is presented with a sequence of randomly ordered stimuli and then asked to reproduce the sequence in order, after a short delay. These tasks generally involve the visual or auditory presentation of letters, numbers, words, or pictures, after which the subject has to verbally repeat the sequence or point to items in the correct order.

For TD individuals, serial recall tasks are thought to recruit primarily verbal rehearsal processes in working memory (i.e. *phonological* verbal representations), for instance as evidenced by decreased memory spans for long words—the *word length effect*—or for phonologically similar items—the *phonological similarity effect* (Baddeley, 2003). These verbal effects are seen even with visually presented stimuli in TD children above seven years of age, suggesting that in later development, TD individuals tend to recode visual stimuli into a verbal form (Hitch et al. 1989). In younger TD children, there is evidence for visual (and not verbal) encoding of visual stimuli in the form of decreased memory spans for visually similar items—the *visual similarity effect* (Hitch et al. 1989b).

Several published studies on serial recall tasks show no significant group differences in overall performance between individuals with autism and controls.² As with the *n*-back task, these data are often used to indicate intact verbal working memory in autism. For example, standardized tests such as the WISC and the WRAML use number and letter span subtests as components of verbal IQ, and individuals with autism have often shown peaks of ability on these particular subtests (Siegel et al. 1996). However, additional behavioral data, such as the presence or absence of the word length or similarity effects described in the previous paragraph, should be considered to determine what strategy an individual is actually using.

Two studies have examined the robustness of the word length effect in individuals with autism. Russell et al. (1996) found, for auditorily presented stimuli, no difference in word length effect in a verbal response condition between children with autism and TD controls as well as a group with moderate learning disabilities, but, oddly, the autism group's word length effect actually increased in a nonverbal (pointing) response condition. In contrast, Whitehouse et al. (2006) used visually presented stimuli with verbal responses and found a smaller word length effect in the autism group than in TD controls. Also, the word length effect increased in the autism group in an overt labeling condition, suggesting that the autism group may have relied to a lesser extent on verbal encoding than controls when not biased to do so by having to produce labels.

Williams et al. (2008) looked at a similar recall task with visually presented stimuli and verbal responses and measured the robustness of the phonological similarity and visual similarity effects in children with autism and in a control group with learning disabilities. They found no group differences in recall performance, but when subjects were divided by their verbal mental

² See Appendix B.

age (VMA), those with VMA over 7 years had better overall recall performance and a significant phonological similarity effect but no visual similarity effect, while subjects with VMA less than 7 years exhibited the opposite pattern. In other words, this study found VMA to better predict strategy use than did diagnostic group, and additional analyses found VMA to be a better predictor than cognitive profiles as well (Williams & Jarrod, 2010). While the authors of this study did not discount the significance of cognitive profile in predicting strategy use, they cautioned against treating it as the only variable of relevance, and they also pointed out the importance of looking at variables like VMA and cognitive profile, in addition to diagnostic group, in assessing results in experimental studies of autism. On both of these points, we wholeheartedly agree, and the question of how to experimentally identify and analyze data from subgroups within the ASD population is central to the continued development of the TiP hypothesis.

In summary, many studies have reported individuals with autism achieving similar levels of performance on serial recall tasks as TD individuals, but at least some of these studies have found evidence of a visual strategy bias in autism.

Dual Task Studies

Dual task studies aim to discern task strategy choices by looking at whether executing a simultaneous secondary task interferes with performance (Brooks, 1968). The basic assumption of the dual-task paradigm is that, because different cognitive modalities (e.g. visual versus verbal) draw upon separate and limited cognitive resources, performing two tasks simultaneously using the same modality will degrade performance more than performing two tasks that use different modalities (Jonides et al. 1996; Navon & Gopher, 1979). Whether a *primary task* uses a certain modality can be determined by finding out whether the simultaneous execution of a secondary task known to involve those resources affects performance (Baddeley & Hitch, 1974). Secondary tasks (a.k.a. *suppression tasks*) can be very simple, so there is little ambiguity about what cognitive resources are being used. Verbal or articulatory suppression (i.e. recruiting *phonological* verbal representations) often consists of repeating a word out loud. Visuospatial suppression can include holding an image in memory or performing a simple tapping or pointing task.

Dual task studies offer a good test of the TiP hypothesis, because their results can clearly indicate, for a particular individual or group, whether visual or verbal cognitive resources are necessary for some primary task. In particular, across a range of primary tasks typically done verbally (tasks for which controls show impairments under verbal but not visual suppression), our hypothesis predicts that individuals with autism will show impairments under visual but not verbal suppression. Only a handful of dual task studies have been performed with individuals on the autism spectrum, and although none have had exactly this form, all have shown results generally consistent with the TiP hypothesis, though not necessarily interpreted as such.

García-Villamizar and Della Sala (2002) used a primary task of serial recall, with verbal recall of auditorily presented digits, and a secondary suppression task of visuomotor tracking, in which subjects had to manually mark a series of boxes on paper. No group differences were found for either task performed singly, but when performed together, the autism group showed a significant impairment on both tasks, while the control group showed no impairment. The authors read these results as marking a general deficit in simultaneous task performance in autism, but these data could also indicate that the group with autism was using a visual strategy

for the digit span task, which, unlike the verbal strategy used by controls, was open to interference from the visual suppression task. Moreover, as discussed below, other dual task studies in autism have not found evidence of a general dual-tasking deficit.

Whitehouse et al. (2006) conducted a dual-task experiment in which the primary task was task-switching in written arithmetic, in which subjects had to alternately add and subtract pairs of numbers, and the secondary task was verbal suppression, with subjects repeating “Monday” out loud. No group differences were found in latency or accuracy in the single-task condition. However, the control group showed an increase in latency under articulatory suppression, matching previous studies on task switching in TD individuals (Baddeley et al. 2001; Emerson & Miyake, 2003), while the autism group did not. These results go against the idea of a general impairment in dual task performance in autism and also suggest that the autism group used a nonverbal (though not necessarily visual) task-switching strategy. Lidstone et al. (2009) re-analyzed these data divided by cognitive profile and found that the lack of a latency increase under articulatory suppression was limited to children with autism having a $V < NV$ profile, irrespective of absolute levels of verbal ability. Controls with a $V < NV$ profile did show impaired dual task performance under articulatory suppression, as did children with a $V = NV$ profile in both groups. Wallace et al. (2009) looked at the Tower of London planning task as the primary task, with a secondary task of articulatory suppression, and similarly found that the control group showed a significant impairment in their primary task performance under articulatory suppression, whereas the autism group showed no such impairment.

Holland and Low (2010) repeated the task switching experiment of Whitehouse et al. (2006) but with an added visuospatial suppression task, with subjects tapping out a simple pattern on a set of blocks using their non-dominant hand. As in the study by Whitehouse et al. (2006), there were no significant group differences in latency or accuracy in the single-task condition. Dual task results showed that the autism group exhibited an increase in task-switching latency under visuospatial suppression but not under articulatory suppression, while the control group showed a similar latency increase under both suppression conditions. Similar dual-task results were obtained in a second experiment that looked at a Tower of Hanoi planning task. At first glance, these data seem to suggest that the autism group used visuospatial but not verbal resources for task-switching and planning, while controls used both visuospatial and verbal resources for both tasks. However, in the task-switching experiment, both groups also showed an increase in latency under visuospatial suppression for a baseline, non-task-switching version of the arithmetic task, suggesting that the visuospatial suppression task may have interfered with peripheral, non-task-switching demands of the primary task. For instance, the visuomotor demands of tapping blocks with the non-dominant hand while writing arithmetic answers with the dominant hand may have been in contention, in which case the visuospatial suppression task did not really target high-level task-switching resources.

While none of these dual-task studies taken singly provides a definitive test of the TiP hypothesis, together they are highly suggestive of individuals with autism using visual strategies for certain tasks that are typically done verbally.

Raven’s Progressive Matrices

Raven’s Progressive Matrices (RPM) is a standardized intelligence test that consists of problems resembling geometric analogies, in which a matrix of figures is presented with one entry missing and the correct missing entry must be selected from among a set of answer choices (Raven,

1936). Although the test is only supposed to measure *eductive* ability, or the ability to extract and understand information from a complex situation, factor analyses have shown the RPM to be a good measure of Spearman's *g*, and it is thus widely used as a general intelligence test (Raven et al. 2003). Using the RPM as a measure of general intelligence, though it consists only of problems in a single format, stands in contrast to using broader tests like the Wechsler scales, which contain subtests across several different domains.

Whereas the RPM scores of TD individuals are usually correlated with their Wechsler IQ scores, individuals with autism have demonstrated RPM scores much higher than their Wechsler scores (Bölte et al. 2009; Dawson et al. 2007; Mottron, 2004). Individuals with Asperger's have shown a similar pattern (Hayashi et al. 2008). One possible explanation for these results is that the RPM, in which both questions and answers are presented visually, might be amenable to solution using visual strategies. The Wechsler scales, on the other hand, are heavily verbal, and while individuals with autism often show good performance on certain subtests like Digit Span or Block Design, their performance on the other subtests can be much lower. In contrast, if TD individuals draw from a combination of visual and verbal strategies on both types of tests, then we would expect to see correlations in their scores between the two paradigms.

One widely cited computational modeling study proposes that TD individuals use a propositional, rule-based strategy to solve RPM problems (Carpenter et al. 1990). However, Hunt (1974) proposed the existence of two qualitatively distinct strategies: one visual, using perceptual operations like visual continuity and superposition, and one analytic, using formal operations based on logical rules. Several behavioral studies of TD individuals point to the possibility of distinct visual and verbal strategies being effective on the RPM (DeShon et al. 1995; Lynn et al. 2004; van der Ven & Ellis, 2000), and we are currently conducting computational studies to investigate whether visual-only algorithms can, in fact, successfully solve the RPM (Kunda et al. 2010a, 2010b).

Soulières et al. (2009) recently found, using fMRI, that individuals with autism had lower brain activation in verbal prefrontal and parietal areas and higher activation in visual occipital areas than TD controls while solving the RPM, consistent with the notion of a visual-strategy-bias in autism. On a related but non-RPM set of matrix reasoning tasks, Sahyoun et al. (2009) found evidence, through measures of response latency, of the autism group having a bias towards visuospatial mediation, whereas TD individuals and individuals with Asperger's were able to use verbal mediation.

Semantic Processing

Evidence from neuropsychology has suggested that visual and verbal semantic memory are somewhat dissociated, in that brain lesions can selectively impair the use of one or the other (Hart & Gordon, 1992). However, whether this dissociation reflects two separate, modality-specific semantic stores or a single store with multiple, modality-specific access schemes is unclear (Caramazza, 1996; Farah & McClelland, 1991). Either way, under the TiP hypothesis, we predict that individuals with autism have privileged or primary access to visual semantic information, whereas TD individuals are capable of accessing both visual and verbal semantics.

In one well-designed fMRI study, Kana et al. (2006) studied brain activation in individuals with autism and TD individuals while they answered true/false questions about high or low imagery sentences. High imagery sentences included statements like, "The number eight when rotated 90 degrees looks like a pair of eyeglasses," while low imagery sentences included

statements like, “Addition, subtraction, and multiplication are all math skills.” One way to conceptualize these two classes of stimuli is as follows:

- (a) High imagery sentences require semantic understanding plus visual reasoning.
- (b) Low imagery sentences require semantic understanding only.

The control group showed a significant difference between the high and low imagery conditions, with the high imagery condition eliciting more activity from temporal and parietal regions associated with mental imagery as well as from inferior frontal regions associated with verbal processing. This pattern fits the model that visual regions are used for visual reasoning, while verbal regions are used for lexical and semantic processing. (The baseline used for both conditions was a fixation task that involved no linguistic processing.) In contrast, the autism group showed similar activation in both conditions, with less activity in inferior frontal language regions than the control group in the high imagery condition, and greater activity in occipital and parietal visual regions in the low imagery condition. This pattern suggests that the individuals with autism may have used visual regions for both visual reasoning and semantic processing.

Many other studies have found significant differences in brain activity during semantic processing tasks between individuals with autism and TD controls, although the precise patterns of results have varied. Like the study by Kana et al. (2006), Gaffrey et al. (2007) found increased activation in posterior visual regions and decreased activation in frontal verbal regions for individuals with ASD during a task of determining whether a word belonged to certain semantic categories (tools, colors, and feelings), with a baseline perceptual processing task. However, Just et al. (2004), in a study of sentence comprehension with a fixation baseline, found reduced activity in visual, occipito-parietal regions in subjects with autism compared to TD controls, though the autism group did also show decreased activity in frontal language regions. Harris et al. (2006) found similar results of reduced frontal language region activation in an ASD group compared to TD controls during a word judging task with a perceptual processing baseline, and also found that the ASD group showed more similar activation in some language regions between the semantic and perceptual tasks than did the control group. In contrast, Knaus et al. (2008) used a response-naming task with a perceptual processing baseline and found that subjects with ASD had greater activation in frontal and temporal language areas than did TD controls.

One important factor in neuroimaging studies of semantic processing is the choice of a baseline task. For TD individuals, lexical-semantic tasks are often paired with perceptual processing tasks that use letter or word stimuli, in order to remove any perceptual components of the semantic understanding process. However, if a subject uses visual neural machinery to do semantic processing, then it is possible that subtracting the brain activation due to a perceptual processing task may remove semantic-related activation in visual regions as well.

In addition to these neuroimaging studies, several behavioral studies have also looked at semantic processing in individuals with autism. Kamio and Toichi (2000) used a word-completion task in which semantic priming was provided using either picture cues or word cues. TD controls performed similarly under both conditions, but the autism group performed much better with picture cues than word cues, suggesting that they were better able to retrieve verbal information through pictorial representations than through other verbal representations. Lopez and Leekam (2003) found that children with autism were as capable as TD controls of using visual semantic context to facilitate object identification; the same pattern was found for verbal semantic information, though ceiling effects were a possible confound in the verbal case.

In summary, while existing data are mixed, current modality-specific models of semantic memory (whether modality-specific in indexing alone or in storage as well) make semantic processing a good candidate for further testing of the TiP hypothesis.

False Belief Tasks

False belief tasks represent one experimental paradigm for testing theory of mind abilities, which center on the attribution of mentalistic or belief states to external entities. Theory of mind, in turn, represents one component of social cognition. False belief tasks comprise one domain that is widely found to be impaired among individuals on the autism spectrum (see review in Happé, 1995), and deficits in theory of mind (e.g. Mindblindness) and other aspects of social cognition have been suggested to be a central facet of autism (Baron-Cohen, 1995; Baron-Cohen & Belmonte, 2005).

One classic test of false belief understanding is the Sally-Anne task (Wimmer & Perner, 1983), in which the subject is shown a skit with two dolls, Sally and Anne. Sally places a marble into a basket and, after Anne leaves the room, moves the marble from the basket into a box. The subject is then asked where Anne will look for the marble when she returns. Responding correctly, that Anne will look in the basket, requires an understanding of Anne's *false belief* that the marble is still in the basket; Anne's belief is false in that it represents something that the subject watching the skit knows is not true.

Many interpretations of false belief task performance in autism posit that there is some fundamentally social deficit that leads to impaired theory of mind abilities (e.g. Baron-Cohen, 1995). We investigate one contrasting view, namely that false belief impairments in autism stem from a domain-general bias against using verbal representations, not from a domain-specific difference in social cognition. In particular, verbal mental age has been found to be strongly correlated with performance on false belief tasks in both individuals with autism and in TD controls (Happé, 1995; Yirmiya et al. 1998). While this pattern seems amenable to a straightforward TiP interpretation, it raises the question of precisely how verbal mental representations might be related to false belief tasks.

One possibility is that standard false belief tasks, which require explicit language comprehension and responding, overtax the weak language skills of individuals with autism. However, individuals with autism also show impairments on nonverbal analogues of false-belief tasks such as eye-tracking studies, making this explanation unlikely (Senju et al. 2009; Senju et al. 2010).

A second possibility is that *linguistic* verbal mental representations are required for developing concepts of false belief, on which both verbal and nonverbal versions of false-belief tasks rely (e.g. Fernyhough, 2008). However, two-year-old TD infants exhibit visual attentional patterns that seem to draw upon an understanding of false beliefs before significant linguistic abilities have developed (Southgate et al. 2007). While there is almost certainly a strong connection between linguistic representations and theory of mind abilities, these types of eye-tracking studies cast doubt on whether the relationship is strictly causal and sequential.

A third possibility, which we espouse, is that verbal representations are, after all, used to form false belief concepts, but where "verbal" in this case refers to *propositional* representations, not *linguistic* representations. Propositions can be thought of as the building blocks of a low-level representational system, where a single proposition takes the form of a related set of symbols that carries semantic meaning. Linguistic representations occur at a much higher level of abstraction than propositions and are explicitly tied to a particular language.

The idea of false belief impairments in autism having a low-level representational origin is not new; constructing false belief concepts has been described as requiring, for instance, the representation of complements (de Villiers & de Villiers, 2003; Hale & Tager-Flusberg, 2003) or meta-representation (Leslie, 1987). The gist of these arguments is that, in order to represent a false belief, an individual must have some mechanism for representing a statement as being held to be true in one context (e.g. as believed by an agent in a story), alongside the property of its being false in a different context (e.g. in the story itself). Recent modeling work in cognitive architectures has found that this type of information structure can be easily represented using propositions (Bello & Cassimatis, 2006).

From this perspective, individual performance on “mental” and “non-mental” versions of false belief tasks should be correlated. While for a time, several visual tasks such as the false photograph, false map, and false drawing tasks were thought to be appropriate non-mental analogues of false belief tasks (e.g. Leekam & Perner, 1991; Charman & Baron-Cohen, 1992; Leslie & Thaiss, 1992), Perner and Leekam (2008) have argued that these tasks do not tap the same representational structure as standard false belief tasks. Instead, they propose that the false sign (or false signal) task is the more appropriate non-mental analogue, and in support of their claim, correlated patterns of impairments have been observed in autism on the false signal task and standard false belief tasks (Bowler et al. 2005). These results support the view of false belief competency being more a function of domain-general *representational* ability than of domain-specific *social* ability.

If false belief impairments in autism are due to deficits in underlying propositional representations, then false belief tasks may seem to fall under the first TiP prediction, regarding tasks only solvable verbally. However, there have been some recent attempts to help individuals with autism represent false belief concepts visually, for instance using thought bubbles or photograph-in-the-head analogies (McGregor et al. 1998a, 1998b; Swettenham et al. 1996; Wellman et al. 2002). These studies have generally shown positive results in teaching subjects to pass specific false belief tasks but less success in leading subjects to transfer their knowledge to new tasks.

TiP Prediction #2: Tasks Typically Done Visually

Unlike the tasks discussed previously, which are typically done verbally but might be amenable to visual strategies, we now discuss empirical data related to the second TiP prediction, on tasks that are done visually both by TD individuals and by individuals with autism. As described earlier, while the TiP hypothesis provides a good explanation of why individuals with autism might show intact performance on these types of tasks, it does not provide a straightforward explanation of superior performance, and it is inconsistent with evidence of performance decrements. In this section, we discuss data that fall into both of these categories.

Visual Search

One widely reported area of superior performance for individuals on the autism spectrum is visual search. For example, individuals on the spectrum have repeatedly demonstrated more accurate and/or more efficient performance on the Embedded Figures Task (EFT), in which a small figure must be located within a larger, more complex one (see review in Happé & Frith, 2006). Several recent papers have looked at classic target/distracter visual search tasks and have

found similar patterns of superior performance by individuals on the autism spectrum, often through faster response latencies.³ Moreover, faster search performance in autism often grows more pronounced with more difficult search tasks, e.g. for conjunctive vs. feature search, etc.

Studies of the EFT using fMRI have shown that individuals with autism tend to recruit more occipital visual processing brain regions for this task, whereas TD controls recruit more frontal and parietal working memory regions (Manjaly et al. 2007; Ring et al. 1999). However, looking at a target/distracter search task, Keehn et al. (2008) found increased activation in individuals on the autism spectrum compared to TD controls in both frontoparietal and occipital regions. This study also found that, while patterns of activation differed for controls between an easy feature search task and a more difficult one, no such differences were found for the autism group. In addition, significant group differences in eye-movement patterns (Keehn et al. 2009) and in sensitivity to task parameters (Baldassi et al. 2009) have been found on visual search tasks. These results are often explained by theories that posit processing strengths in autism, and in particular, some recent evidence suggests that enhanced low-level perceptual discrimination may contribute to faster search in autism (Joseph et al. 2009).

In general, many studies point to the existence of significant and widespread differences between individuals on the autism spectrum and TD individuals on visual search tasks and in overall patterns of visual attention, and these differences seem to developmentally precede many other cognitive processes (Brenner et al. 2007). Specific relationships between the TiP hypothesis and visual search and attention remain to be determined, especially in terms of development and basic perceptual processes.

Spatial Recall

Serial spatial recall tasks are a part of many standardized intelligence tests, such as Finger Windows in the WRAML. These tasks involve the presentation of a sequence of spatial locations (e.g. holes on a card or blocks on a table), which the subject has to manually reproduce. Another type of spatial recall task uses self-ordered pointing, in which the subject must point to locations not previously selected. Both paradigms require the subject to reproduce a set or sequence of spatial locations. Individuals with autism often, but not always, show impaired performance on these types of tasks, and we found no study of spatial recall on which the autism group showed superior performance.⁴

Given that serial recall for items or objects appears to be unimpaired in autism, as discussed earlier, there appears to be a dissociation between how well individuals with autism can remember visually discriminable items vs. visually indiscriminable spatial locations. Although these results seem to contradict the TiP hypothesis, one explanation could be that the visual representations used by individuals with autism do not, by themselves, represent spatial information adequately. In line with this idea, on tasks that combine visual and spatial information (i.e. recalling the locations of visually discriminable stimuli), individuals with autism have shown intact performance (Ozonoff & Strayer, 2001; Williams et al. 2006).

Another possibility might be that spatial recall tasks actually recruit verbal working memory; correlations between spatial span and speech rate have been found in TD individuals,

³ See Appendix C.

⁴ See Appendix D.

without similar correlations between spatial span and tapping or spatial movement rate (Chuah & Maybery, 1999; Smyth & Scholey, 1992, 1996). Studies have also found that articulatory suppression can interfere with spatial span tasks (Jones et al. 1995; Smyth et al. 1988; Smyth & Pelky, 1992).

Visual Recall

One paradigm for tests of visual recall involves giving the subject an abstract design to draw from memory after an initial inspection. Two examples are the Benton Visual Retention Test and the Rey-Osterrieth Complex Figure Task. The Rey-Osterrieth task includes a copy condition that helps to identify perceptual or motor impairments that could confound results.

Many studies of these types of tasks have revealed decreased performance in individuals with autism.⁵ Given the patterns of intact and even superior performance found in other visual domains, these visual recall data are rather puzzling. Moreover, both the Rey and Benton tests have been found, in TD individuals, to be correlated with the Block Design subtest of the Wechsler scales and not correlated with verbal measures (Mitrushina et al. 2005; Strauss et al. 2006), and the Block Design subtest has been commonly cited as an area of particular strength for individuals with autism (Siegel et al. 1996).

One explanation could be that perceptual and motor components of these drawing tasks are what cause difficulties for individuals with autism rather than the memory requirements per se. Ropar and Mitchell (2001) examined this possibility by comparing differences in copy and recall scores among experimental groups, instead of just looking at recall scores alone, and found no group differences between TD controls and subjects with autism or Asperger's. Alternately, individuals with autism could have difficulty on the spatial but not visual aspects of these tasks. Although the Rey-Osterrieth task is often described as a test of visual memory, the task contains both visual and spatial components that are somewhat dissociable (Breier et al. 1996).

As with spatial recall, data on visual recall for individuals with autism are mixed at best. It is unclear how these results might be accounted for by the TiP hypothesis, and more detailed investigations are needed of what specific cognitive processes both individuals with autism and TD individuals recruit for these tasks.

Discussion

We have presented detailed reviews of empirical data on individuals with autism from several different task domains. For each task, we have attempted to give an objective assessment of whether the data are consistent with our formulation of a Thinking in Pictures (TiP) hypothesis about cognition in autism. As expected, the results of this analysis are mixed. Certain task domains offer evidence that is highly consistent with and well explained by the TiP hypothesis, including: (1) the n-back task, (2) serial recall, (3) dual tasking, (4) Raven's Progressive Matrices, (5) semantic processing, and (6) false belief tasks. Other task domains, while not inconsistent with the TiP hypothesis, are not directly explained by it either, namely: (7) visual

⁵ See Appendix E.

search. Finally, there are task domains whose data seem to contradict the TiP hypothesis, which are: (8) spatial recall, and (9) visual recall.

Of course, there are many experimental task paradigms that we have not addressed or have only briefly touched upon, for instance free recall, cued recall, visual or verbal recognition, executive functioning, etc. However, the main point that we wish to convey is that, across several task domains, there is a significant amount of evidence that is highly consistent with the TiP hypothesis. This finding is even more interesting given that most of the studies we reviewed did not explicitly use a visual/verbal hypothesis in the design or execution of their experiments.

In the remainder of this section, we discuss the relationship of the TiP hypothesis to existing cognitive accounts of autism. We close with our final thoughts on TiP and important questions for further exploration.

Theories of Cognition in Autism

Several existing cognitive theories of autism aim to explain various aspects of autistic behavior. We briefly discuss three of these theories here—Executive Dysfunction, Weak Central Coherence, and Enhanced Perceptual Functioning—focusing on how the TiP hypothesis relates to them, on points of congruence as well as divergence. (Possible relationships of TiP to a fourth theory, Mindblindness, were covered in the section describing false belief tasks.)

The Executive Dysfunction (ED) theory posits that autism is characterized by impairments in a set of higher-level cognitive skills that underlie independent, goal-oriented behavior, such as planning, set-shifting, and generativity (Russell, 1997). We argue that evidence in support of the ED theory is consistent with the TiP hypothesis if the specific executive capacities found to be impaired in autism are those that cannot be performed using visual mental representations. For example, individuals with autism are often impaired on the Wisconsin Card Sorting Test (WCST), a test of set-shifting in which subjects must maintain knowledge of a sorting rule and then switch the rule as needed (see review in Hill, 2004). The WCST, however, has been found to rely heavily on language abilities and verbal working memory in TD individuals (Baldo et al. 2005). More generally, Russell et al. (1999) propose that individuals with autism may have trouble primarily with executive tasks that require the implicit verbal encoding of rules. However, despite these suggestive pieces of data, evaluating a potential link between executive functioning in autism and the TiP hypothesis will require a close re-examination of a wide range of tasks used to tap executive abilities to discern how they fit into the task decomposition presented earlier (i.e. can they be solved visually, verbally, or using either type of mental representation).

The Weak Central Coherence (WCC) theory suggests that individuals with autism may exhibit a bias towards local over global processing (Happé & Frith, 2006). Much of the evidence for the WCC theory shows patterns of either poor performance in individuals with autism on tasks that are said to rely on global processing of stimuli, or intact or superior performance on tasks that are said to rely on local processing. However, at least some of the “local” tasks cited by the WCC theory are visual, e.g. embedded figures, block design, visual search, etc. Likewise, certain WCC “global” tasks are verbal, e.g. homograph pronunciation. For at least these tasks, the TiP hypothesis can provide an explanation that is consistent with published data, although, as mentioned earlier, the TiP hypothesis does not currently provide a concrete explanation of autistic superiorities on certain tasks, beyond our speculation that a reliance on visual representations might lead to increased visual expertise. Moreover, the WCC literature has

identified several non-visual local tasks that are also performed well by individuals with autism, such as pitch and melody perception (see review in Happé & Frith, 2006). The TiP hypothesis is, at present, silent about representational modalities other than visual or verbal, though these results raise the question of whether TiP can (or should) be extended to a more general perceptual/verbal distinction.

Along these lines, the Enhanced Perceptual Functioning (EPF) theory proposes that individuals with autism have enhanced low level perceptual processing across a variety of modalities, in contrast to cognitive processing that involves higher levels of neural integration (Mottron et al. 2006). For instance, several studies have found evidence of atypicalities, and often superiorities, in low-level visual perception in autism (e.g. Bertone et al. 2005; Vandembroucke et al. 2008). In addition to low-level perceptual enhancements and atypicalities, Ropar and Mitchell (2002) have proposed that autistic perception can be characterized, at least in certain task domains, as being less influenced than in TD individuals by “top-down” cognitive processes that draw upon prior conceptual knowledge. Caron et al. (2006) suggest that a combination of locally oriented processing and enhanced perceptual processing leads to superiorities in autism on visual tasks, for the subgroup of individuals who share these two traits.

Unlike the TiP hypothesis, which at present focuses only on visual representations, EPF and other perceptual accounts of autism are stated broadly to encompass a variety of perceptual modalities. However, within consideration of the visual modality, there seems to be significant overlap between these accounts, especially in that both TiP and EPF propose “a successful, problem-solving use of perceptual [brain] areas” (Mottron et al. 2006). Also, inasmuch as working with verbal representations might fall under “high-level” cognition, additional overlaps between TiP and EPF are likely.

One major difference between the WCC and EPF theories and the TiP hypothesis is that WCC and EPF embody *process* accounts of cognition, equating various representational modalities—visual, auditory, etc.—within each of two distinct types of processing—local vs. global, or perceptual vs. high-level. TiP, on the other hand, embodies a *content* account of cognition, equating various processing types—perception, working memory, long-term memory, etc.—within each of two distinct representational modalities—visual vs. verbal. Another difference is that WCC and EPF more explicitly account for autistic superiorities on certain visual tasks, whereas TiP does not currently propose a concrete mechanism for this pattern of performance, though several possibilities, such as increased visual expertise, remain to be explored. It is plausible that these accounts are linked, both developmentally and cognitively, and the precise relationship between the TiP hypothesis and these theories is the subject of some of our current work.

Final Thoughts

Our results lead us to propose two main conclusions. First, given the existence of considerable evidence in line with the TiP hypothesis, the idea that certain individuals with autism may “think visually” should be taken seriously as a cognitive model and receive more focused and sustained attention in behavioral and neurobiological experiments. Second, and more generally, the interpretation of behavioral data from individuals with autism (or, indeed, with any form of atypical cognition) should be performed with care. Assumptions governing the relations between cognition and behavior that hold for TD individuals may not hold universally, and we

have presented several instances in which visual and verbal strategies seem to be recruited differently across experimental groups, despite often producing superficially similar behavior.

If a subset of individuals on the autism spectrum does have a bias towards using visual mental representations, then several important questions remain to be answered about the TiP hypothesis. How might this subset of individuals be identified, and how could experimental subgroups be appropriately defined to account for cognitive differences within the autism spectrum? Would these individuals display a $V < NV$ profile, and would such a profile be a necessary and/or sufficient marker of their cognitive style? What, if anything, might the TiP hypothesis tell us about individuals on the spectrum who showed $V = NV$ or $V > NV$ cognitive profiles? At present, we do not have answers to these questions.

Other important avenues for further inquiry include (1) the accuracy and interpretation of measures of visual and verbal IQ when potential differences in task strategies are taken into account, (2) the distinction, if any, between visual and spatial processing under the TiP account, as well as relationships with other types of perceptual processing, (3) how biases away from using verbal representations and towards using visual representations might be causally linked, and (4) what role TiP might play in neurobiological and developmental accounts of autism.

Author Note

Author information: Maithilee Kunda and Ashok K. Goel, Design & Intelligence Lab, School of Interactive Computing, Georgia Institute of Technology.

This research was supported in part by a grant (IIS Award 0534266) on “Multimodal Case-Based Reasoning in Modeling and Design” from the National Science Foundation, by a National Defense Science and Engineering Graduate Fellowship from the Office of Naval Research, and by a Graduate Research Fellowship from the National Science Foundation. The authors would like to thank A. Rozga for her careful reading of this manuscript, K. McGregor and E. Schumacher for many discussions on this topic, and G. Abowd, R. Arriaga, and the rest of the Autism Research Group at the Georgia Tech/Emory Health Systems Institute for their encouragement and support of this work.

Correspondence concerning this article should be addressed to Maithilee Kunda, School of Interactive Computing, Georgia Institute of Technology, 85 Fifth Street NW, Atlanta, GA 30332. Email: mkunda@gatech.edu

References

- Ambery, F., Russell, A., Perry, K., Morris, R., & Murphy, D. (2006). Neuropsychological functioning in adults with Asperger syndrome. *Autism, 10*, 551-564.
- Ameli, R., Courchesne, E., Lincoln, A., Kaufman, A., & Grillon, C. (1988). Visual memory processes in high-functioning individuals with autism. *Journal of Autism and Developmental Disorders, 18*, 601-615.
- American Psychiatric Association. (2000). *Diagnostic and Statistical Manual of Mental Disorders* (Revised 4th Ed.). Washington, DC: American Psychiatric Association.
- Baddeley, A. (2003). Working memory: Looking back and looking forward. *Nature Rev. Neuroscience, 4*, 829-839.
- Baddeley, A., Chincotta, D., & Adlam, A. (2001). Working memory and the control of action: Evidence from task switching. *Journal of Experimental Psychology: General, 130*, 641-657.
- Baddeley, A., & Hitch, G. (1974). Working memory. In G. Bower (Ed.), *The Psychology of Learning and Motivation* (Vol. 8, pp. 47-89). New York: Academic Press.
- Baldassi, S., Pei, F., Megna, N., Recupero, G., Viespoli, M., Iglizzi, R., Tancredi, R., et al. (2009). Search superiority in autism within, but not outside the crowding regime. *Vision Research, 49*, 2151-2156.
- Baldo, J., Dronkers, N., Wilkins, D., Ludy, C., Raskin, P., & Kim, J. (2005). Is problem solving dependent on language? *Brain and Language, 92*, 240-250.
- Baron-Cohen, S. (1995). *Mindblindness: An essay on autism and theory of mind*. Cambridge, MA: MIT Press.
- Baron-Cohen, S., & Belmonte, M. (2005). Autism: A window onto the development of the social and the analytic brain. *Annual Review of Neuroscience, 28*, 109-126.
- Bello, P., & Cassimatis, N. (2006). Developmental accounts of theory-of-mind acquisition: Achieving clarity via computational cognitive modeling. In *Proc. 28th Annual Conf. Cognitive Science Society*, pp. 1014-1019.
- Bennetto, L., Pennington, B., & Rogers, S. (1996). Intact and impaired memory functions in autism. *Child Development, 67*, 1816-1835.
- Bertone, A., Mottron, L., Jelenic, P., & Faubert, J. (2005). Enhanced and diminished visuo-spatial information processing in autism depends on stimulus complexity. *Brain, 128*, 2430-2441.
- Bölte, S., Dziobek, I., & Poustka, F. (2009). Brief report: The level and nature of autistic intelligence revisited. *Journal of Autism and Developmental Disorders, 39*, 678-682.
- Bowler, D., Briskman, J., Gurvidi, N., & Fornells-Ambrojo, M. (2005). Understanding the mind or predicting signal-dependent action? Performance of children with and without autism on analogues of the false-belief task. *Journal of Cognition and Development, 6*, 259-283.
- Breier, J., Plenger, P., Castillo, R., Fuchs, K., Wheless, J., Thomas, A., et al. (1996). Effects of temporal lobe epilepsy on spatial and figural aspects of memory for a complex geometric figure. *Journal of the International Neuropsychological Society, 2*, 535-540.
- Brenner, L., Turner, K., & Müller, R. (2007). Eye movement and visual search: Are there elementary abnormalities in autism? *Journal of Autism and Developmental Disorders, 37*, 1289-1309.
- Brooks, L. (1968). Spatial and verbal components of the act of recall. *Canadian Journal of Psychology, 22*, 349-368.
- Caramazza, A. (1996). Pictures, words, and the brain. *Nature, 383*, 216-217.
- Caron, M., Mottron, L., Rainville, C., & Chouinard, S. (2004). Do high functioning persons with autism present superior spatial abilities? *Neuropsychologia, 42*, 467-481.
- Caron, M., Mottron, L., Berthiaume, C., & Dawson, M. (2006). Cognitive mechanisms, specificity and neural underpinnings of visuospatial peaks in autism. *Brain, 129*, 1789-1802.
- Carpenter, P., Just, M., & Shell, P. (1990). What one intelligence test measures: A theoretical account of the processing in the Raven Progressive Matrices Test. *Psychological Review, 97*, 404-31.
- Charman, T., & Baron-Cohen, S. (1992). Understanding drawings and beliefs: A further test of the metarepresentation theory of autism: A research note. *J. Child Psychology and Psychiatry, 33*, 1105-1112.
- Chuah, Y., & Maybery, M. (1999). Verbal and spatial short-term memory: Common sources of developmental change? *Journal of Experimental Child Psychology, 73*, 7-44.
- Dawson, M., Soulières, I., Gernsbacher, M., & Mottron, L. (2007). The level and nature of autistic intelligence. *Psychological Science, 18*, 657-662.
- DeShon, R., Chan, D., & Weissbein, D. (1995). Verbal overshadowing effects on Raven's Advanced Progressive Matrices: Evidence for multidimensional performance determinants. *Intelligence, 21*, 135-155.
- de Villiers, J., & de Villiers, P. (2003). Language for thought: Coming to understand false beliefs. In *Language in Mind: Advances in the Study of Language and Thought* (pp. 335-384). Cambridge, MA: MIT Press.

- Edgin, J., & Pennington, B. (2005). Spatial cognition in autism spectrum disorders: Superior, impaired, or just intact? *Journal of Autism and Developmental Disorders*, *35*, 729-745.
- Emerson, M., & Miyake, A. (2003). The role of inner speech in task switching: A dual-task investigation. *Journal of Memory and Language*, *48*, 148-168.
- Farah, M., & McClelland, J. (1991). A computational model of semantic memory impairment: Modality specificity and emergent category specificity. *Journal of Experimental Psychology: General*, *120*, 339-357.
- Fernyhough, C. (2008). Getting Vygotskian about theory of mind: Mediation, dialogue, and the development of social understanding. *Developmental Review*, *28*, 225-262.
- Gaffrey, M., Kleinhans, N., Haist, F., Akshoomoff, N., Campbell, A., Courchesne, E., et al. (2007). Atypical participation of visual cortex during word processing in autism: An fMRI study of semantic decision. *Neuropsychologia*, *45*, 1672-1684.
- García-Villamizar, D., & Della Sala, S. (2002). Dual-task performance in adults with autism. *Cognitive Neuropsychiatry*, *7*, 63-74.
- Grandin, T. (2006). *Thinking in pictures, expanded edition: My life with autism*. New York, NY: Vintage Books.
- Gunter, H., Ghaziuddin, M., & Ellis, H. (2002). Asperger syndrome: Tests of right hemisphere functioning and interhemispheric communication. *Journal of Autism and Developmental Disorders*, *32*, 263-281.
- Hale, C., & Tager-Flusberg, H. (2003). The influence of language on theory of mind: A training study. *Developmental Science*, *6*, 346-359.
- Happé, F. (1995). The role of age and verbal ability in the theory of mind task performance of subjects with autism. *Child Development*, *66*, 843-855.
- Happé, F., & Frith, U. (2006). The weak coherence account: Detail-focused cognitive style in autism spectrum disorders. *Journal of Autism and Developmental Disorders*, *36*, 5-25.
- Harris, G., Chabris, C., Clark, J., Urban, T., Aharon, I., Steele, S., et al. (2006). Brain activation during semantic processing in autism spectrum disorders via functional magnetic resonance imaging. *Brain and Cognition*, *61*, 54-68.
- Hart, J., & Gordon, B. (1992). Neural subsystems for object knowledge. *Nature*, *359*, 60-64.
- Hayashi, M., Kato, M., Igarashi, K., & Kashima, H. (2008). Superior fluid intelligence in children with Asperger's disorder. *Brain and Cognition*, *66*, 306-310.
- Heaton, P., Ludlow, A., & Roberson, D. (2008). When less is more: Poor discrimination but good colour memory in autism. *Research in Autism Spectrum Disorders*, *2*, 147-156.
- Hill, E. (2004). Evaluating the theory of executive dysfunction in autism. *Developmental Review*, *24*, 189-233.
- Hitch, G., Halliday, M., Dodd, A., & Littler, J. (1989). Development of rehearsal in short-term memory: Differences between pictorial and spoken stimuli. *British Journal of Developmental Psychology*, *7*, 347-362.
- Hitch, G., Woodin, M., & Baker, S. (1989b). Visual and phonological components of working memory in children. *Memory and Cognition*, *17*, 175-185.
- Holland, L., & Low, J. (2010). Do children with autism use inner speech and visuospatial resources for the service of executive control? Evidence from suppression in dual tasks. *British Journal of Developmental Psychology*, *28*, 369-391.
- Hunt, E. (1974). Quote the raven? Nevermore! In L. Gregg (Ed.), *Knowledge and cognition* (pp. 129-158). Hillsdale, NJ: Erlbaum.
- Hurlburt, R., Happé, F., & Frith, U. (1994). Sampling the form of inner experience in three adults with Asperger syndrome. *Psychological Medicine*, *24*, 385-396.
- Jarrold, C., Gilchrist, I., & Bender, A. (2005). Embedded figures detection in autism and typical development: Preliminary evidence of a double dissociation in relationships with visual search. *Developmental Science*, *8*, 344-351.
- Jolliffe, T., & Baron-Cohen, S. (1997). Are people with autism and Asperger syndrome faster than normal on the embedded figures test? *Journal of Child Psychology and Psychiatry and Allied Disciplines*, *38*, 527-534.
- Jones, D., Farrand, P., Stuart, G., & Morris, N. (1995). Functional equivalence of verbal and spatial information in serial short-term memory. *J. Experimental Psychology: Learning, Memory, and Cognition*, *21*, 1008-1018.
- Jonides, J., Reuter-Lorenz, P., Smith, E., Awh, E., Barnes, L., Drain, M., et al. (1996). Verbal and spatial working memory in humans. *Psychology of learning and motivation*, *35*, 43-88.
- Joseph, R., Keehn, B., Connolly, C., Wolfe, J., & Horowitz, T. (2009). Why is visual search superior in autism spectrum disorder? *Developmental Science*, *12*, 1083-1096.
- Joseph, R., Steele, S., Meyer, E., & Tager-Flusberg, H. (2005). Self-ordered pointing in children with autism: Failure to use verbal mediation in the service of working memory? *Neuropsychologia*, *43*, 1400-1411.

- Joseph, R., Tager-Flusberg, H., & Lord, C. (2002). Cognitive profiles and social-communicative functioning in children with autism spectrum disorder. *Journal of Child Psychology and Psychiatry*, *43*, 807-821.
- Just, M., Cherkassky, V., Keller, T., & Minshew, N. (2004). Cortical activation and synchronization during sentence comprehension in high-functioning autism: Evidence of underconnectivity. *Brain*, *127*, 1811-1821.
- Kamio, Y., & Toichi, M. (2000). Dual access to semantics in autism: Is pictorial access superior to verbal access? *Journal of Child Psychology and Psychiatry and Allied Disciplines*, *41*, 859-867.
- Kana, R., Keller, T., Cherkassky, V., Minshew, N., & Just, M. (2006). Sentence comprehension in autism: Thinking in pictures with decreased functional connectivity. *Brain*, *129*, 2484-2493.
- Keehn, B., Brenner, L., Palmer, E., Lincoln, A., & Müller, R. (2008). Functional brain organization for visual search in ASD. *Journal of the International Neuropsychological Society*, *14*, 990-1003.
- Keehn, B., Brenner, L., Ramos, A., Lincoln, A., Marshall, S., & Müller, R. (2009). Brief report: Eye-movement patterns during an Embedded Figures test in children with ASD. *Journal of Autism and Developmental Disorders*, *39*, 383-387.
- Kirchner, W. (1958). Age differences in short-term retention of rapidly changing information. *Journal of Experimental Psychology*, *55*, 352-358.
- Klin, A., Volkmar, F., Sparrow, S., Cicchetti, D., & Rourke, B. (1995). Validity and neuropsychological characterization of Asperger syndrome: Convergence with nonverbal learning disabilities syndrome. *Journal of Child Psychology and Psychiatry*, *36*, 1127-1140.
- Knaus, T., Silver, A., Lindgren, K., Hadjikhani, N., & Tager-Flusberg, H. (2008). fMRI activation during a language task in adolescents with ASD. *Journal of the International Neuropsychological Society*, *14*, 967-979.
- Koshino, H., Carpenter, P., Minshew, N., Cherkassky, V., Keller, T., & Just, M. (2005). Functional connectivity in an fMRI working memory task in high-functioning autism. *NeuroImage*, *24*, 810-821.
- Koshino, H., Kana, R., Keller, T., Cherkassky, V., Minshew, N., & Just, M. (2008). fMRI investigation of working memory for faces in autism: Visual coding and underconnectivity with frontal areas. *Cerebral Cortex*, *18*, 289-300.
- Kosslyn, S., Thompson, W., & Ganis, G. (2006). *The case for mental imagery*. New York, NY: Oxford University Press.
- Kunda, M., & Goel, A. (2008). How thinking in pictures can explain many characteristic behaviors of autism. In *Proceedings of the 7th IEEE International Conference on Development and Learning* (pp. 304-309).
- Kunda, M., McGreggor, K., & Goel, A. (2010a). Can the Raven's Progressive Matrices intelligence test be solved by thinking in pictures? Oral presentation given at *IMFAR-2010*, Philadelphia, PA.
- Kunda, M., McGreggor, K., & Goel, A. (2010b). Taking a look (literally!) at the Raven's intelligence test: Two visual solution strategies. In *Proceedings of the 32nd Annual Conference of the Cognitive Science Society*.
- Leekam, S., & Perner, J. (1991). Does the autistic child have a metarepresentational deficit? *Cognition*, *40*, 203-218.
- Leslie, A. (1987). Pretense and representation: The origins of "theory of mind." *Psychological Review*, *94*, 412-426.
- Leslie, A., & Thaiss, L. (1992). Domain specificity in conceptual development: Neuropsychological evidence from autism. *Cognition*, *43*, 225-251.
- Lidstone, J., Fernyhough, C., Meins, E., & Whitehouse, A. (2009). Brief report: Inner speech impairment in children with autism is associated with greater nonverbal than verbal skills. *Journal of Autism and Developmental Disorders*, *39*, 1222-1225.
- Lincoln, A., Courchesne, E., Kilman, B., Elmasian, R., & Allen, M. (1988). A study of intellectual abilities in high-functioning people with autism. *Journal of Autism and Developmental Disorders*, *18*, 505-524.
- Lopez, B., & Leekam, S. (2003). Do children with autism fail to process information in context? *Journal of Child Psychology and Psychiatry*, *44*, 285-300.
- Luna, B., Minshew, N., Garver, K., Lazar, N., Thulborn, K., Eddy, W., et al. (2002). Neocortical system abnormalities in autism: An fMRI study of spatial working memory. *Neurology*, *59*, 834-840.
- Lynn, R., Allik, J., & Irwing, P. (2004). Sex differences on three factors identified in Raven's Standard Progressive Matrices. *Intelligence*, *32*, 411-424.
- Manjaly, Z., Bruning, N., Neufang, S., Stephan, K., Brieber, S., Marshall, J., et al. (2007). Neurophysiological correlates of relatively enhanced local visual search in autistic adolescents. *NeuroImage*, *35*, 283-291.
- McGregor, E., Whiten, A., & Blackburn, P. (1998a). Teaching theory of mind by highlighting intention and illustrating thoughts: A comparison of their effectiveness with 3-year-olds and autistic individuals. *British Journal of Developmental Psychology*, *16*, 281-300.
- McGregor, E., Whiten, A., & Blackburn, P. (1998b). Transfer of the picture-in-the-head analogy to natural contexts to aid false belief understanding in autism. *Autism*, *2*, 367-387.

- Minshew, N., & Goldstein, G. (2001). The pattern of intact and impaired memory functions in autism. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, *42*, 1095-1101.
- Minshew, N., Goldstein, G., Muenz, L., & Payton, J. (1992). Neuropsychological functioning in nonmentally retarded autistic individuals. *Journal of Clinical and Experimental Neuropsychology*, *14*, 749-61.
- Minshew, N., Goldstein, G., & Siegel, D. (1997). Neuropsychologic functioning in autism: Profile of a complex information processing disorder. *Journal of the International Neuropsychological Society*, *3*, 303-316.
- Minshew, N., Luna, B., & Sweeney, J. (1999). Oculomotor evidence for neocortical systems but not cerebellar dysfunction in autism. *Neurology*, *52*, 917-922.
- Mitrushina, M., Boone, K., & Razani, J. (2005). *Handbook of normative data for neuropsychological assessment*. New York: Oxford University Press.
- Morris, R., Rowe, A., Fox, N., Feigenbaum, J., Miotto, E., & Howlin, P. (1999). Spatial working memory in Asperger's syndrome and in patients with focal frontal and temporal lobe lesions. *Brain & Cognition*, *41*, 9-26.
- Mottron, L. (2004). Matching strategies in cognitive research with individuals with high-functioning autism: Current practices, instrument biases, and recommendations. *J. Autism and Developmental Disorders*, *34*, 19-27.
- Mottron, L., Dawson, M., Soulières, I., Hubert, B., & Burack, J. (2006). Enhanced perceptual functioning in autism: An update, and eight principles of autistic perception. *J. Autism and Developmental Disorders*, *36*, 27-43.
- Navon, D., & Gopher, D. (1979). On the economy of the human processing system. *Psychological Review*, *86*, 214-255.
- O'Connor, N., & Hermelin, B. (1967). Auditory and visual memory in autistic and normal children. *Journal of Intellectual Disability Research*, *11*, 126-131.
- O'Riordan, M. (2000). Superior modulation of activation levels of stimulus representations does not underlie superior discrimination in autism. *Cognition*, *77*, 81-96.
- O'Riordan, M. (2004). Superior visual search in adults with autism. *Autism*, *8*, 229-248.
- O'Riordan, M., & Plaisted, K. (2001). Enhanced discrimination in autism. *Quarterly Journal of Experimental Psychology A*, *54*, 961-979.
- O'Riordan, M., Plaisted, K., Driver, J., & Baron-Cohen, S. (2001). Superior visual search in autism. *Journal of Experimental Psychology: Human Perception and Performance*, *27*, 719-730.
- Ozonoff, S., & Strayer, D. (2001). Further evidence of intact working memory in autism. *Journal of Autism and Developmental Disorders*, *31*, 257-263.
- Perner, J., & Leekam, S. (2008). The curious incident of the photo that was accused of being false: Issues of domain specificity in development, autism, and brain imaging. *Quarterly J. Experimental Psychology*, *61*, 76-89.
- Plaisted, K., O'Riordan, M., & Baron-Cohen, S. (1998). Enhanced visual search for a conjunctive target in autism: A research note. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, *39*, 777-783.
- Pylyshyn, Z. (2002). Mental imagery: In search of a theory. *Behavioral and Brain Sciences*, *25*, 157-182.
- Raven, J., Raven, J. C., & Court, J. (2003). *Manual for Raven's Progressive Matrices and Vocabulary Scales, Section I: General overview*. San Antonio, TX: Harcourt Assessment.
- Raven, J. C. (1936). *Mental tests used in genetic studies: The performances of related individuals in tests mainly educative and mainly reproductive* (Unpublished master's thesis). University of London, London.
- Ring, H., Baron-Cohen, S., Wheelwright, S., Williams, S., Brammer, M., Andrew, C., et al. (1999). Cerebral correlates of preserved cognitive skills in autism: A functional MRI study of embedded figures task performance. *Brain*, *122*, 1305-1315.
- Ropar, D., & Mitchell, P. (2001). Susceptibility to illusions and performance on visuospatial tasks in individuals with autism. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, *42*, 539-549.
- Ropar, D., & Mitchell, P. (2002). Shape constancy in autism: The role of prior knowledge and perspective cues. *Journal of Child Psychology & Psychiatry & Allied Disciplines*, *43*, 647-653.
- Russell, J. (1997). *Autism as an executive disorder*. New York: Oxford University Press.
- Russell, J., Jarrold, C., & Henry, L. (1996). Working memory in children with autism and with moderate learning difficulties. *Journal of Child Psychology and Psychiatry*, *37*, 673-686.
- Russell, J., Jarrold, C., & Hood, B. (1999). Two intact executive capacities in children with autism: Implications for the core executive dysfunctions in the disorder. *J. Autism and Developmental Disorders*, *29*, 103-112.
- Sahyoun, C., Soulières, I., Belliveau, J., Mottron, L., & Mody, M. (2009). Cognitive differences in pictorial reasoning between high-functioning autism and Asperger's syndrome. *Journal of Autism and Developmental Disorders*, *39*, 1014-1023.
- Senju, A., Southgate, V., Miura, Y., Matsui, T., Hasegawa, T., Tojo, Y., Osanai, H., et al. (2010). Absence of spontaneous action anticipation by false belief attribution in children with autism spectrum disorder. *Development and Psychopathology*, *22*, 353-360.

- Senju, A., Southgate, V., White, S., & Frith, U. (2009). Mindblind eyes: An absence of spontaneous theory of mind in Asperger Syndrome. *Science*, *325*, 883-885.
- Siegel, D., Minshew, N., & Goldstein, G. (1996). Wechsler IQ profiles in diagnosis of high-functioning autism. *Journal of Autism and Developmental Disorders*, *26*, 389-406.
- Smith, E., & Jonides, J. (1999). Storage and executive processes in the frontal lobes. *Science*, *283*, 1657-1661.
- Smyth, M., Pearson, N., & Pendleton, L. (1988). Movement and working memory: Patterns and positions in space. *Quarterly Journal of Experimental Psychology A*, *40*, 497-514.
- Smyth, M., & Pelky, P. (1992). Short-term retention of spatial information. *British J. Psychology*, *83*, 359-374.
- Smyth, M., & Scholey, K. (1992). Determining spatial span: The role of movement time and articulation rate. *Quarterly Journal of Experimental Psychology A*, *45*, 479-501.
- Smyth, M., & Scholey, K. (1996). The relationship between articulation time and memory performance in verbal and visuospatial tasks. *British Journal of Psychology*, *87*, 179-191.
- Soulières, I., Dawson, M., Samson, F., Barbeau, E., Sahyoun, C., Strangman, G., et al. (2009). Enhanced visual processing contributes to matrix reasoning in autism. *Human Brain Mapping*, *30*, 4082-4107.
- Southgate, V., Senju, A., & Csibra, G. (2007). Action anticipation through attribution of false belief by 2-year-olds. *Psychological Science*, *18*, 587-592.
- Steele, S., Minshew, N., Luna, B., & Sweeney, J. (2007). Spatial working memory deficits in autism. *Journal of Autism and Developmental Disorders*, *37*, 605-612.
- Strauss, E., Sherman, E., & Spreen, O. (2006). *A compendium of neuropsychological tests: Administration, norms, and commentary*. New York: Oxford University Press.
- Swettenham, J., Baron-Cohen, S., Gomez, J., & Walsh, S. (1996). What's inside someone's head? Conceiving of the mind as a camera helps children with autism acquire an alternative to a theory of mind. *Cognitive Neuropsychiatry*, *1*, 73-88.
- Vandenbroucke, M., Scholte, H., van Engeland, H., Lamme, V., & Kemner, C. (2008). A neural substrate for atypical low-level visual processing in autism spectrum disorder. *Brain*, *131*, 1013-1024.
- van der Ven, A. & Ellis, J. (2000). A Rasch analysis of Raven's Standard Progressive Matrices. *Personality and Individual Differences*, *29*, 45-64.
- Verté, S., Geurts, H., Roeyers, H., Oosterlaan, J., & Sergeant, J. (2005). Executive functioning in children with autism and Tourette syndrome. *Development and Psychopathology*, *17*, 415-445.
- Verté, S., Geurts, H., Roeyers, H., Oosterlaan, J., & Sergeant, J. (2006). Executive functioning in children with an Autism Spectrum Disorder: Can we differentiate within the spectrum? *Journal of Autism and Developmental Disorders*, *36*, 351-372.
- Wallace, G., Silvers, J., Martin, A., & Kenworthy, L. (2009). Brief report: Further evidence for inner speech deficits in autism spectrum disorders. *Journal of Autism and Developmental Disorders*, *39*, 1735-1739.
- Wellman, H., Baron-Cohen, S., Caswell, R., Gomez, J., Swettenham, J., Toye, E., et al. (2002). Thought-bubbles help children with autism acquire an alternative to a theory of mind. *Autism*, *6*, 343-363.
- Whitehouse, A., Maybery, M., & Durkin, K. (2006). Inner speech impairments in autism. *Journal of Child Psychology and Psychiatry*, *47*, 857-865.
- Williams, D., Happe, F., & Jarrold, C. (2008). Intact inner speech use in autism spectrum disorder: Evidence from a short-term memory task. *Journal of Child Psychology and Psychiatry*, *49*, 51-58.
- Williams, D., Goldstein, G., Carpenter, P., & Minshew, N. (2005). Verbal and spatial working memory in autism. *Journal of Autism and Developmental Disorders*, *35*, 747-756.
- Williams, D., Goldstein, G., & Minshew, N. (2006). The profile of memory function in children with autism. *Neuropsychology*, *20*, 21-29.
- Williams, D., & Jarrold, C. (2010). Brief report: Predicting inner speech use amongst children with autism spectrum disorder (ASD): The roles of verbal ability and cognitive profile. *Journal of Autism and Developmental Disorders*, *40*, 907-913.
- Wimmer, H., & Perner, J. (1983). Beliefs about beliefs: Representation and constraining function of wrong beliefs in young children's understanding of deception. *Cognition*, *13*, 103-128.
- Yirmiya, N., Erel, O., Shaked, M., & Solomonica-Levi, D. (1998). Meta-analyses comparing theory of mind abilities of individuals with autism, individuals with mental retardation, and normally developing individuals. *Psychological Bulletin*, *124*, 283-307.

Appendix A

Review of Results from Published Studies of the n-Back Task in Autism

Reference	N	Age	VIQ	PIQ	FSIQ	Task details	Results
Koshino et al. 2005	14	25.7 (10.2)	102.6 (14.1)	—	100.1	$n = 0, 1, 2$; visually presented letters.	NSGD on accuracy or RT. Significant group differences in patterns of brain activation.
Koshino et al. 2008	11	24.5 (10.2)	106.1 (14.1)	102.1 (13.8)	104.5 (13.1)	$n = 0, 1, 2$; grayscale face pictures.	NSGD on accuracy or RT. Significant group differences in patterns of brain activation.
Ozonoff & Strayer, 2001	25	12.94 (3.18)	94.6 (18.5)	99.3 (19.9)	96.3 (17.8)	$n = 1, 2$; visually presented colored shapes.	NSGD on accuracy or RT. RT tended to be correlated with VIQ.
Williams et al. 2005	31	26.58 (8.68)	111.10 (16.47)	103.13 (16.64)	108.65 (16.75)	$n = 0, 1, 2$; visually presented letters.	NSGD on accuracy or RT.
	24	11.75 (2.36)	112.50 (16.53)	106.38 (14.21)	109.67 (16.07)	$n = 0, 1, 2$; visually presented letters.	NSGD on accuracy or RT.

Note. N = number of participants; VIQ, PIQ, and FSIQ = verbal, performance, and full-scale IQ, respectively; NSGD = no significant group differences; RT = reaction time. Age and IQ values are shown as: *mean (standard deviation)* in years and scores, respectively. All subjects diagnosed with autism, all control groups TD, and NSGD in age or shown IQ measures.

Appendix B

Review of Results from Published Studies of Serial Recall Tasks in Autism

Reference	N	Age	VIQ	PIQ	FSIQ	Items	Presentation	Response	Results
Ameli et al. 1988	16 ^f	22.7 (4.9)	81 (16)	90.6 (13.5)	83 (14)	digits	auditory	verbal	NSGD in digit span.
Bennetto et al. 1996	19 ^{ac}	15.95 (3.3)	82.32 (15.2)	98.11 (15.9)	88.89 (11.1)	digits	auditory	verbal	NSGD in digit span.
Joseph et al. 2005	24 ^a	8.9 (2.3)	94 (19)	99 (20)	96 (18)	words pictures	auditory visual	pointing self-ordered pointing	NSGD in correct responses. NSGD in errors in nonverbal condition. AU had more errors in verbal than nonverbal condition and than controls.
Minshew et al. 1992	15	21.13 (8.02)	98.53 (21.63)	92.87 (10.72)	95.73 (13.61)	digits	auditory	verbal	NSGD in digit span.
Minshew et al. 1997	33	20.91 (9.69)	102.48 (16.35)	97.45 (11.19)	100.09 (12.96)	digits	auditory	verbal	NSGD in correct responses.
Minshew & Goldstein, 2001	52	22.33 (9.59)	94.96 (17.56)	91.52 (12.95)	92.88 (15.06)	letters words	auditory auditory	verbal verbal	NSGD in correct sequences. AU impaired in number of correct sequences.
O'Connor & Hermelin, 1967	12 ^b	11.8	---	---	---	words	auditory	verbal	NSGD in number recalled. Greater position and recency effects in AU than controls.
Ozonoff & Strayer, 2001	25 ^e	12.94 (3.18)	94.6 (18.5)	99.3 (19.9)	96.3 (17.8)	pictures colored boxes	visual visual	manual ordering self-ordered pointing	NSGD in number recalled. NSGD in number of perseverative errors, which tended to correlation with VIQ.

Russell et al. 1996	33 ^{dh}	12.38 (2.95)	---	---	---	words	auditory	verbal, pointing	MLD had lower span than other groups. AU had greater nonverbal WLE than verbal WLE and than MLD group.
Whitehouse et al. 2006	23 ⁱ	11	---	---	---	pictures	visual	verbal	NSGD in correct responses. AU had smaller WLE than controls and in silent vs. label condition.
Williams et al. 2005	24	11.75 (2.36)	112.50 (16.53)	106.38 (14.21)	109.67 (16.07)	digits, letters	auditory	verbal	NSGD in score. Correlated with FSIQ in AU but not in controls.
Williams et al. 2006	38	11.68 (2.46)	106.42 (15.97)	100.55 (14.19)	103.82 (14.29)	digits, letters	auditory	verbal	NSGD in score.
Williams et al. 2008	25 ^{bc}	12.25 (3.08)	77.16 (15.25)	76.84 (20.27)	74.84 (15.99)	pictures	visual	verbal	NSGD in span. Subjects with high VMA had greater PSE than VSE; vice versa for low VMA.

Note. N = number of participants; VIQ, PIQ, and FSIQ = verbal, performance, and full-scale IQ, respectively; NSGD = no significant group differences; AU = autism group; MLD = mild learning disabilities group; WLE, PSE, and VSE = word length, phonological similarity, and visual similarity effects, respectively; VMA = verbal mental age. Age and IQ values are shown as: *mean (standard deviation)* in years and scores, respectively. All subjects diagnosed with autism, all control groups TD, and NSGD in age or shown IQ measures, except as noted.

^a Subject group diagnosed as autism/PDD-NOS. ^b Subject group diagnosed as ASD. ^c Control group comprised of individuals with moderate learning difficulties (MLD). ^d Controls grouped as TD and MLD. ^e Controls grouped as TD and Tourette's Syndrome. ^f Subject group had slightly lower PIQ than controls, but NSGD in age. ^g NSGD in nonverbal ability according to Peabody Picture Vocabulary Test. ^h NSGD in VMA, but AU had higher age. ⁱ NSGD in verbal, nonverbal, and reading ability, but AU had higher age.

Appendix C

Review of Results from Published Studies of Target/Distracter Visual Search Tasks in Autism

Reference	N	Age	CPM	Search	Target	Distracters	Accuracy	RT
Jarrod et al. 2005	18 ^b	12.42 (1.98)	21.56 (7.45)	F	red jumping clown	green skinny clown, red fat clown	---	AU faster.
				C	red jumping clown	green jumping clown, red skinny clown	---	AU faster.
Keehn et al. 2008	9 ^a	15.1 (2.5)	---	F	upright <i>T</i>	right-pointing <i>T</i>	NSGD.	NSGD.
				F	upright <i>T</i>	right, left, bottom-pointing <i>T</i>	NSGD.	NSGD.
O'Riordan, 2000	11	9.2 (0.8)	27 (4)	C	red <i>X</i>	red <i>T</i> , green <i>X</i>	NSGD.	AU faster.
				C	red <i>X</i>	red letters and colored <i>X</i>	NSGD.	AU faster.
				C	red <i>X</i>	red <i>T</i> , green <i>X</i>	NSGD.	AU faster.
				C	green <i>T</i>	red <i>T</i> , green <i>X</i>	NSGD.	AU faster.
				C	red <i>X</i> , green <i>T</i>	red <i>T</i> , green <i>X</i>	NSGD.	AU faster.
O'Riordan, 2004	10	22.0 (3.6)	---	F	<i>N</i>	<i>P</i> , <i>Q</i>	NSGD.	NSGD.
				C	<i>R</i>	<i>P</i> , <i>Q</i>	NSGD.	AU faster.
				F	ellipse	circle	NSGD.	AU faster.
				C	red <i>X</i>	green <i>X</i> , red <i>C</i>	NSGD.	NSGD.
				C	red <i>F</i>	pink <i>F</i> , red <i>E</i>	AU more accurate.	NSGD.
O'Riordan & Plaisted, 2001	15	9.2 (1.1)	28 (3)	C	vertical red bar	vertical green bar, horizontal red bar	NSGD.	AU faster.
				C	vertical red bar	vertical green line, horizontal green bar, horizontal green line	NSGD.	NSGD.
				C	vertical red bar	vertical green bar, horizontal red bar, vertical red line	NSGD.	AU faster.
				C	red <i>X</i>	green <i>X</i> , red <i>C</i>	NSGD.	NSGD.
				C	red <i>X</i>	red <i>X</i>	NSGD.	NSGD.

	C	red X	pink X, red C	NSGD.	AU faster.
	C	red F	green F, red E	NSGD.	AU faster.
	C	red F	pink F, red E	NSGD.	AU faster.
	F	red S	green X, red T	NSGD.	NSGD.
O'Riordan et al. 2001	12	8.4 (0.9)	26 (4)	NSGD.	AU faster.
	F	tilted line	vertical lines	NSGD.	NSGD.
	F	vertical line	tilted lines	NSGD.	AU faster.
Plaisted et al. 1998	8 ^c	8.8 (1.2)	---	NSGD.	NSGD.
	C	red X	green X, red T	AU more accurate.	AU faster.

Note. N = number of participants; CPM = Raven's Colored Progressive Matrices; RT = reaction time; F = feature search; C = conjunctive search; AU = autism group; NSGD = no significant group differences. Age and CPM values are shown as: *mean (standard deviation)* in years and raw scores, respectively. All subjects diagnosed with autism, all control groups TD, and NSGD in age or cognitive measures, except as noted.

^a Subject group diagnosed as ASD. ^b AU had higher chronological age. ^c Autism group had higher block design scores.

Appendix D

Review of Results from Published Studies of Spatial Recall Tasks in Autism

Reference	N	Age	VIQ	PIQ	FSIQ	Task	Details	Results
Caron et al. 2004	16 ^a	17.6 (6.3)	102.2 (21.2)	112.3 (12.9)	107.7 (13.1)	Maze route learning	learn choice points in an actual maze	NSGD in accuracy or speed of going through maze.
Edgin & Pennington, 2005	24 ^a	11.46 (2.32)	104.40 (20.24)	---	---	CANTAB Spatial Working Memory	touch one of identical boxes to find tokens (self-ordered pointing)	NSGD in errors or strategy. Both correlated with age and block design score in both groups.
Luna et al. 2002	11	32.3 (9.3)	106.7 (14.9)	96.5 (10.5)	102.7 (12.1)	Oculomotor Delayed Response	shift gaze to previous stimulus location	AU impaired in saccade accuracy compared to controls. Not impaired on baseline saccade task.
Minshew & Goldstein, 2001	52	22.33 (9.59)	94.96 (17.56)	91.52 (12.95)	92.88 (15.06)	Maze recall	learn choice points in hidden maze	AU impaired on complex mazes but not on simpler maze.
Minshew et al. 1992	15	21.13 (8.02)	98.53 (21.63)	92.87 (10.72)	95.73 (13.61)	Maze recall	learn choice points in hidden maze	NSGD.
Minshew et al. 1997	33	20.91 (9.69)	102.48 (16.35)	97.45 (11.19)	100.09 (12.96)	Maze recall	learn choice points in hidden maze	NSGD.
Minshew et al. 1999	26	20.2 (8.5)	98.5 (16.9)	90.1 (12.7)	94.0 (14.1)	Oculomotor Delayed Response	shift gaze to previous stimulus location	AU impaired in saccade accuracy compared to controls. Not impaired on baseline saccade task.
Morris et al. 1999	15 ^b	29.5	99	100.1	---	Executive Golf Task	touch one of identical golf holes to find putts (self-ordered pointing)	AS impaired on within and between search errors.
Steele et al. 2007	29	14.83 (5.47)	107.52 (13.02)	106.21 (11.82)	107.76 (10.99)	CANTAB Spatial Working Memory	touch one of identical boxes to find tokens (self-ordered pointing)	AU showed greater errors and lower strategy score, both correlated with PIQ but not VIQ. TD scores not correlated with any IQ.
Verté et al. 2005	61 ^{de}	9.1 (1.9)	---	---	99.2 (17.1)	Corsi block tapping	forward tapping recall of blocks	AU impaired.
Verté et al. 2006	50 ^f	8.7 (1.9)	93.1 (18.0)	104.0 (17.8)	98.2 (17.3)	Corsi block tapping	forward tapping recall of blocks	AU impaired.

	37 ^b	8.5 (2.1)	105.6 (20.0)	104.0 (15.9)	105.2 (16.3)	Corsi block tapping	forward tapping recall of blocks	AS impaired.
	25 ^{ef}	8.5 (1.4)	93.3 (14.7)	105.8 (19.2)	98.3 (14.4)	Corsi block tapping	forward tapping recall of blocks	NSGD.
Williams et al. 2005	31	26.58 (8.68)	111.10 (16.47)	103.13 (16.64)	108.65 (16.75)	WMS-III spatial span	forward and backward tapping recall of blocks	AU impaired. Correlated with FSIQ in AU but not in controls.
Williams et al. 2006	24	11.75 (2.36)	112.50 (16.53)	106.38 (14.21)	109.67 (16.07)	Finger Windows	poke sequence of holes in a card	AU impaired. No correlation with FSIQ in either group.
Williams et al. 2006	38	11.68 (2.46)	106.42 (15.97)	100.55 (14.19)	103.82 (14.29)	Finger Windows	poke sequence of holes in a card	AU impaired.

Note. N = number of participants; VIQ, PIQ, and FSIQ = verbal, performance, and full-scale IQ, respectively; NSGD = no significant group differences; AU = autism group; AS = Asperger's group. Age and IQ are shown as: *mean (standard deviation)* in years and scores, respectively. All subjects diagnosed with autism, all controls TD, and NSGD in age or shown IQ measures, except as noted.

^a Subject group diagnosed as Autism/Asperger's. ^b Subject group diagnosed as Asperger's. ^c Subject group diagnosed as PDD-NOS.

^d Controls grouped as TD, Tourette's, and autism/Tourette's (comorbid). ^e Subject group had lower FSIQ. ^f Subject group had lower FSIQ and VIQ.

Appendix E

Review of Results from Published Studies of Visual Recall Tasks in Autism

Reference	N	Age	VIQ	PIQ	FSIQ	Task	Details	Results
Ambery et al. 2006	27 ^a	37.6 (14.6)	106.1 (15.7)	103.7 (19.2)	---	Doors and People Test	draw shapes from memory, also recognize colored photos of doors	AU impaired on combined measure of visual recall and recognition.
Ameli et al. 1988	16 ^e	22.7 (4.9)	81 (16)	90.6 (13.5)	83 (14)	BVRT	draw series of abstract figures from memory	AU impaired. Highly correlated with PIQ in AU but not in controls.
Gunter et al. 2002	8 ^a	16.25 (10.19)	111.38 (15.22)	96.13 (13.13)	---	Rey-Osterrieth	draw abstract figure from memory	NSGD in copy/recall scores or score differences.
Minshew & Goldstein, 2001	52	22.33 (9.59)	94.96 (17.56)	91.52 (12.95)	92.88 (15.06)	Rey-Osterrieth	draw abstract figure from memory	AU impaired in number of elements correctly drawn for immediate and delayed recall.
Minshew et al. 1997	15	21.13 (8.02)	98.53 (21.63)	92.87 (10.72)	95.73 (13.61)	Rey-Osterrieth	draw abstract figure from memory	AU impaired in number of elements correctly drawn for delayed recall.
Minshew et al. 1992	33	20.91 (9.69)	102.48 (16.35)	97.45 (11.19)	100.09 (12.96)	Rey-Osterrieth	draw abstract figure from memory	AU impaired in number of elements correctly drawn for delayed recall.
Ropar & Mitchell, 2001	19 ^{ef} 11 ^{acf}	14.2 (2.4)	---	---	---	Rey-Osterrieth	draw abstract figure from memory	NSGD in copy/recall score differences or in strategy.
Verté et al. 2005	61 ^{de}	9.1 (1.9)	---	---	99.2 (17.1)	BVRT	draw series of abstract figures from memory	AU impaired.
Verté et al. 2006	50 ^h 37 ^a	8.7 (1.9) 8.5 (2.1)	93.1 (18.0)	104.0 (17.8)	98.2 (17.3)	BVRT	draw series of abstract figures from memory	AU impaired.
			105.6 (20.0)	104.0 (15.9)	105.2 (16.3)	BVRT	draw series of abstract figures from memory	AU impaired.

25 th	8.5 (1.4)	93.3 (14.7)	105.8 (19.2)	98.3 (14.4)	BVRT	draw series of abstract figures from memory	AU impaired.
Williams et al. 2006	38	11.68 (2.46)	106.42 (15.97)	100.55 (14.19)	103.82 (14.29)	WRAML Design Memory draw geometric shapes from memory	AU impaired.

Note. N = number of participants; VIQ, PIQ, and FSIQ = verbal, performance, and full-scale IQ, respectively; NSGD = no significant group differences; AU = autism group, BVRT = Benton Visual Retention Test. Age and IQ values are shown as: *mean (standard deviation)* in years and scores, respectively. All subjects diagnosed with autism, all control groups TD, and NSGD in age or shown IQ measures, except as noted.

^a Subject group diagnosed as Asperger's. ^b Subject group diagnosed as PDD-NOS. ^c Controls grouped as 8-yr.-old TD, 11-yr.-old TD, and learning disabled. ^d Controls grouped as TD, Tourette's, and autism/Tourette's (comorbid). ^e Subject group had lower PIQ. ^f Subject group had higher age. ^g Subject group had lower FSIQ. ^h Subject group had lower FSIQ and VIQ.